Journal of Pharmaceutical and Biomedical Analysis

Volume 13 1995
LIST OF CONTENTS

Number 1

	v	Editorial
	vii	APQ Newsletter
Full papers		
Zhiling Qin	1	Advances in biopharmaceutical analysis in the People's Republic of China: 1991–1993
U. Hammerling, R. Kroon and L. Sjödin	9	*In vitro* bioassay with enhanced sensitivity for human granulocyte colony-stimulating factor
R.B. Taylor and R.G. Reid	21	Analysis of basic antimalarial drugs by CZE; Part 2. Validation and application to bioanalysis
D.G. Watson, Q. Su, J.M. Midgley, E. Doyle and N.S. Morton	27	Analysis of unconjugated morphine, codeine, normorphine and morphine as glucuronides in small volumes of plasma from children
K.D. Altria, T. Wood, R. Kitscha and A. Roberts-McIntosh	33	Validation of a capillary electrophoresis method for the determination of potassium counter-ion levels in an acidic drug salt
B.A. Dawson and D.B. Black	39	Utilizing nuclear magnetic resonance (NMR) spectroscopy for assessing nadolol racemate composition
J.E. Noroski, D.J. Mayo and M. Moran	45	Determination of the enantiomer of a cholesterol-lowering drug by cyclodextrin-modified micellar electrokinetic chromatography
A.F.M. El Walily, A. El Gindy and A.A.M. Wahbi	53	Spectrophotometric determination of flunarizine dihydrochloride through the formation of charge-transfer complex with iodine
B.R. Simmons, N.K. Jagota and J.T. Stewart	59	A supercritical fluid chromatographic method using packed columns for phenylbutazone and oxyphenbutazone in serum, and for phenylbutazone in a dosage form
C. Anchisi, B. Batetta, F. Sanna, A.M. Fadda, A.M. Maccioni and S. Dessi	65	HDL subfractions as altered in cancer patients
Short Communications		
C.G. Kontoyannis	73	Quantitative determination of $CaCO_3$ and glycine in antacid tablets by Laser Raman Spectroscopy
E. Kwong, L. Baert and S. Béchard	77	Gas–liquid chromatographic quantitation of polyethylene glycol 400 in pharmaceutical preparations
J.E. Noroski and D.J. Mayo	83	Chiral LC of a cholesterol-lowering drug using serum albumin mobile phases

Number 2

Full Papers		
D. Dadgar, P.E. Burnett, M.G. Choc, K. Gallicano and J.W. Hooper	89	Application issues in bioanalytical method validation, sample analysis and data reporting
C. McDonald, P. Mortimer, G. McAllister, P.E. Gardiner, J. Reglinski and W.E. Smith	99	A comparative study of the interaction of Myocrisin with albumin and γ-globulin

G. Pyramides, J.W. Robinson and S.W. Zito — 103 — The combined use of DSC and TGA for the thermal analysis of atenolol tablets

F.E. Blondino and P.R. Byron — 111 — The quantitative determination of aspirin and its degradation products in a model solution aerosol

C.-J. Wang, Z. Tian, V. Reyes and C.-C. Lin — 121 — A specific enzyme immunoassay (EIA) with selective extraction for quantitation of a topical anti-inflammatory agent, SCH 40120, in human plasma

S.C. Sharma, M.B. Evans and S.J. Evans — 129 — The enantiomeric separation of metipranolol and desacetyl-metipranolol on a cellulose tris-3,5-dimethylphenyl-carbamate chiral stationary phase

R. Lozano, F.V. Warren Jr, S. Perlman and J.M. Joseph — 139 — Quantitative analysis of fosinopril sodium by capillary zone electrophoresis and liquid chromatography

K.M. Steele and C.E. Lunte — 149 — Microdialysis sampling coupled to on-line microbore liquid chromatography for pharmacokinetic studies

R.D. Amin, H. Cheng and J.D. Rogers — 155 — Determination of MK-0476 in human plasma by liquid chromatography

G. La Follette, A.L. Jayewardene, A.K. Seneveiratne, E.T. Lin and J.G. Gambertoglio — 159 — Determination of ticarcillin in human plasma and in urine by reversed-phase LC

A.S. Chilton, R.E. Godward and P.F. Carey — 165 — The determination in human plasma of 1-hydroxy-2-naphthoic acid following administration of salmeterol xinafoate

Short Communications

T.A. Walker — 171 — A liquid chromatographic assay for citric acid in over-the-counter carbamide peroxide products using indirect UV detection

W.-N. Wu, J.A. Masucci, L.A. McKown and J.D. Rodgers — 177 — Identification of four plasma metabolites of Etoperidone in dogs

E.O. Fadiran, J. Leslie, M. Fossler and D. Young — 185 — Determination of clozapine and its major metabolites in human serum and rat plasma by liquid chromatography using solid-phase extraction and ultraviolet detection

Number 3

Full Papers

M.M. De Villiers — 191 — Influence of cohesive properties of micronized drug powders on particle size analysis

L. de La Peña, A. Gómez-Hens and D. Pérez-Bendito — 199 — Kinetic determination of nortriptyline in pharmaceutical samples by use of photometric and fluorimetric detection

M.L. Anthony, V.S. Rose, J.K. Nicholson and J.C. Lindon — 205 — Classification of toxin-induced changes in ^{1}H NMR spectra of urine using an artificial neural network

T. Jovanović, B. Stanović and Z. Korićanac — 213 — Spectrophotometric investigation on complex formation of captopril with palladium(II) and its analytical application

B. Morelli — 219 — Determination of ternary mixtures of antibiotics, by ratio-spectra zero-crossing first- and third-derivative spectrophotometry

P.G. Pietta, A. Bruno, P.L. Mauri, C. Gardana, R. Maffei-Facino and M. Carini — 229 — Determination of sunscreen agents by micellar electrokinetic chromatography

M.C. Gardia-Alvarez-Coque, E.F. Simo-Alfonso, G. Ramis-Ramos and J.S. Esteve-Romero — 237 — High-performance micellar liquid chromatography determination of sulphonamides in pharmaceuticals after azodye precolumn derivatization

R.A. Rajewski, D.G. Kosednar, T.A. Matches, O.S. Wong, K. Burchett and K. Thakker — 247 — Stereo-specific analysis of a novel protein kinase C inhibitor

A.J.G. Mank, E.J. Molenaar, C. Gooijer, H. Lingeman, N.H. Velthorst and U.A.Th. Brinkman — 255 — Determination of the anticancer drug metabolite WR 1065 using precolumn derivatization and diode laser induced fluorescence detection

A.J. Humberstone, G.J. Currie, C.J.H. Porter, M.J. Scanlon and W.N. Charman — 265 — A simplified liquid chromatography assay for the quantitation of halofantrine and desbutylhalofantrine in plasma and identification of a degradation product of desbutylhalofantrine formed under alkaline conditions

S. Scalia, A. Renda, G. Ruberto, F. Bonina and E. Menegatti — 273 — Assay of vitamin A palmitate and vitamin E acetate in cosmetic creams and lotions by supercritical fluid extraction and HPLC

I.A. Kaltashov, X. Yu and C. Fenselau — 279 — Simple interface for microbore LC and electrospray ionization mass spectrometry and analysis of melphalan–alkylation sites in metallothionein

D.K. Bempong, I.L. Honigberg and N.M. Meltzer — 285 — Normal phase LC–MS determination of retinoic acid degradation products

D.L. Norwood, D. Prime, B.P. Downey, J. Creasey, S.K. Sethi and P. Haywood — 293 — Analysis of polycyclic aromatic hydrocarbons in metered dose inhaler drug formulations by isotope dilution gas chromatography/mass spectrometry

R. De la Torre, J. Ortuño, M.L. González, M. Farré, J. Camí and J. Segura — 305 — Determination of cocaine and its metabolites in human urine by gas chromatography/mass spectrometry after simultaneous use of cocaine and ethanol

S. Carlsen, M. Hayes and M.L. Powell — 313 — Quantitative determination and pharmacokinetics of a new antipsychotic (CGS 13429A) in human plasma using capillary gas chromatography/mass spectrometry

Short Communications

M.T. Maya, J.P. Pais and J.A. Morais — 319 — A rapid method for the determination of piroxicam in plasma by high-performance liquid chromatography

S.W. Baertschi, D.E. Dorman, L.A. Spangle, M.W. Collins and L.J. Lorenz — 323 — Formation of fluorescent pyrazine derivatives via a novel degradation pathway of the carbacephalosporin loracarbef

R. Herrmann — 329 — Automated HPLC assay for sotalol in human plasma

E. Gaetani, C. Laureri and M. Vitto — 335 — GC–ITD detection and quantitative analysis of Proxazole in cows' plasma and milk

Number 4/5

Papers from the Fifth International Symposium on Pharmaceutical and Biomedical Analysis, 21–24 September 1994, Stockholm, Sweden

xi — Editorial

Full Papers

M. Cichna, P. Markl and J.F.K. Huber — 339 — Determination of true octanol–water partition coefficients by means of solvent generated liquid–liquid chromatography

K.-J. Lindner, P. Hartvig, N. Tyrefors, C. Hedlund and B. Långström — 353 — An automated liquid chromatographic plasma analysis of amino acids used in combination with positron emission tomography (PET) for determination of *in vivo* plasma kinetics

K.-J. Lindner, P. Hartvig, J. Tedroff, A. Ljungström, P. Bjurling and B. Långström — 361 — Liquid chromatographic analysis of brain homogenates and microdialysates for the quantification of L-[β-^{11}C]DOPA and its metabolites for the validation of positron emission tomography studies

L. Ståhle, A. Mian and N. Borg	369	Use of multivariate characterization, design and analysis in assay optimization
G. Zackrisson, G. Östling, B. Skagerberg and T. Anfält	377	ACcelerated Dissolution Rate Analysis (ACDRA) for controlled release drugs. Application to Roxiam®
M.T. Gilbert, I. Barinov-Colligon and J.R. Miksic	385	Cross-validation of bioanalytical methods between laboratories
B.A. Dawson, D.B. Black and G.A. Neville	395	Multinuclear NMR (^{1}H, ^{13}C and ^{19}F) spectroscopic re-examination of the solvolytic behaviour of flurazepam dihydrochloride
F.W. Langkilde and A. Svantesson	409	Identification of celluloses with Fourier-Transform (FT) mid-infrared, FT-Raman and near-infrared spectrometry
S.P. Jacobsson, M. Carlsson, U. Jönsson and G. Nilsson	415	Quantitative determination of Sulfasalazine by near-infrared spectroscopy and multivariate analysis in reflectance mode with a fibre-optic probe
F. González and R. Pous	419	Quality control in manufacturing process by near infrared spectroscopy
Z. Xinrong, W.R.G. Baeyens, G. Van Der Weken, A.C. Calokerinos and K. Nakashima	425	Chemiluminescence analysis of captopril: comparison between luminol and rhodamine B-sensitized cerium(IV) methods
S. Furlanetto, P. Gratteri, S. Pinzauti, R. Leardi, E. Dreassi and G. Santoni	431	Design and optimization of the variables in the adsorptive stripping voltammetric determination of rufloxacin in tablets, human plasma and urine
L. Campanella, Z. Aturki, M.P. Sammartino and M. Tomassetti	439	Aspartate analysis in formulations using a new enzyme sensor
Y. Su, M. Tomassetti, M.P. Sammartino, G. Crescentini and L. Campanella	449	A new salicylate ISFET for the determination of salicylic and acetylsalicylic acid in drugs
M.N.M.P. Alçada, J.L.F.C. Lima and M.C.B.S.M. Montenegro	459	FIA titrations of ephedrine in pharmaceutical formulations with a PVC tetraphenylborate tubular electrode
B. De Castro, J.L.F.C. Lima and S. Reis	465	Potentiometric determination of formation constants of copper(II)/bile acid/peptide in aqueous solution
J.M. Nedeljković, V.M. Vasić, V.V. Vuković and T.S. Jovanović	471	Kinetics of the reaction of S-carboxymethyl-L-cysteine with palladium(II) chloride
K. Tomita, K. Nomura, H. Kondo, K. Nagata and H. Tsubota	477	Stabilized enzymatic reagents for measuring glucose, creatine kinase and γ-glutamyltransferase with thermostable enzymes from a thermophile, *Bacillus stearothermophilus*
Y. Yoshimoto, A. Shibukawa, H. Sasagawa, S. Nitta and T. Nakagawa	483	Michaelis–Menten analysis of immobilized enzyme by affinity capillary electrophoresis
S.H. Hansen, I. Bjørnsdottir and J. Tjørnelund	489	Separation of basic drug substances of very similar structure using micellar electrokinetic chromatography
I. Bechet, M. Fillet, Ph. Hubert and J. Crommen	497	Quantitative analysis of non-steroidal anti-inflammatory drugs by capillary zone electrophoresis
M.G. Quaglia, A. Farina, E. Bossu and C. Dellaquila	505	Analysis of non-benzodiazepinic anxiolytic agents by capillary zone electrophoresis
C. Bory, C. Chantin and D. Bozon	511	Capillary electrophoretic analysis of DNA restriction fragments and PCR products for polymorphism and mutation studies in cystic fibrosis and Gaucher's disease
K. Hettiarachchi, Y. Ha, T. Tran and A.P. Cheung	515	Application of HPLC and CZE to the analysis of polyoxometalates

E. Forgács	525	Use of principal component analysis for the evaluation of the retention behaviour of monoamine oxidase inhibitory drugs on β-cyclodextrin column
T. Cserháti, E. Forgács and J. Holló	533	Interaction of taxol and other anticancer drugs with α-cyclodextrin
H. Lamparczyk and P.K. Zarzycki	543	Effect of temperature on separation of estradiol stereoisomers and equilin by liquid chromatography using mobile phases modified with β-cyclodextrin
B.-M. Eriksson and A. Wallin	551	Evaluation of the liquid-chromatographic resolution of indenoindolic racemic compounds on three protein-based chiral stationary phases
S. Klick	563	Evaluation of different injection techniques in the gas chromatographic determination of thermolabile trace impurities in a drug substance
M. Erickson, K.-E. Karlsson, B. Lamm, S. Larrson, L.A. Svensson and J. Vessman	567	Identification of a new by-product detected in metoprolol tartrate
A.S. Kord, I. Tumanova and W.L. Matier	575	A novel HPLC method for determination of EDTA in a cataract inhibiting ophthalmic drug
P. Leroy, M. Barbaras, J.-L. Colin and A. Nicolas	581	Ion-exchange liquid chromatography method with indirect UV detection for the assay of choline in pharmaceutical preparations
R. Gatti, V. Andrisano, A.M. Di Pietra and V. Cavrini	589	Analysis of aliphatic dicarboxylic acids in pharmaceuticals and cosmetics by liquid chromatography (HPLC) with fluorescence detection
V. Andrisano, D. Bonazzi and V. Cavrini	597	HPLC analysis of liquorice triterpenoids — applications to the quality control of pharmaceuticals
L. Iavarone, M. Scandola, F. Pugnaghi and P. Grossi	607	Qualitative analysis of potential metabolites and degradation products of a new antiinfective drug in rat urine, using HPLC with radiochemical detection and HPLC–mass spectrometry
A. Rudolphi, S. Vielhauer, K.-S. Boos, D. Seidel, I.-M. Bäthge and H. Berger	615	Coupled-column liquid chromatographic analysis of epirubicin and metabolites in biological material and its application to optimization of liver cancer therapy
D. Fraier, E. Frigerio, E. Pianezzola, M. Strolin Benedetti, J. Cassidy and P. Vasey	625	A sensitive procedure for the quantitation of free and N-(2 hydroxypropyl)methacrylamide polymer-bound doxorubicin and doxorubicin (PK1) and some of its metabolites, 13-dihydrodoxorubicin, 13-dihydrodoxorubicinone and doxorubicinone, in human plasma and urine by reversed-phase HPLC with fluorimetric detection
J. Knoeller, R. Vogt-Schenkel and M.A. Brett	635	A simple and robust HPLC method for the determination of paroxetine in human plasma
A. Andersson and H. Ehrsson	639	Stability of cisplatin and its monohydrated complex in blood, plasma and ultrafiltrate — implications for quantitative analysis
G.H. Ragab, H. Nohta, M. Kai, Y. Ohkura and K. Zaitsu	645	1,2-Diarylethylenediamines as sensitive pre-column derivatizing reagents for chemiluminescence detection of catecholamines in HPLC
A. Helander, C. Löwenmo and O. Beck	651	Determination of 5-hydroxytryptophol in urine by high-performance liquid chromatography: application of a new post-column derivatization method with fluorometric detection
A. Jankowski, A. Skorek, K. Krzysko, P.K. Zarzycki, R.J. Ochocka and H. Lamparczyk	655	Captopril: determination in blood and pharmacokinetics after single oral dose
J. Liliemark, F. Albertioni, C. Edlund and G. Juliusson	661	Bioavailability and bacterial degradation of rectally administered 2-chloro-2′-deoxyadenosine

Short Communications

R. Andersin and M. Mesilaakso 667 Structure elucidation of 6-chloro-2-methyl-4(1H)-quinazolinone, a photodecomposition product of midazolam

N. Murayama, M. Nakaoka, H. Nomura and H. Hakusui 671 Radioimmunoassay for DS-4574, an anti-allergic agent: development, evaluation and application to human plasma samples

S. Vladimirov, Z. Fiser, D. Agbaba and D. Zivanov-Stakic 675 Spectrophotometric determination of fusidic acid and sodium fusidate in dosage forms

T. Jovanović, Z. Korićanac and B. Stanković 679 Silver electrode in direct potentiometric determination of propylthiouracil in pharmaceutical dosage form

K. Nikolic, M. Bogavac and B. Stanković 683 Coulometric determination of nizatidine

I. Bjørnsdottir and S.H. Hansen 687 Determination of opium alkaloids in opium by capillary electrophoresis

B. Chankvetadze, L. Chankvetadze, Sh. Sidamonidze, E. Yashima and Y. Okamoto 695 Enantioseparation of some chiral pharmaceuticals using narrow-bore liquid chromatography

N. Sadlej-Sosnowska 701 Inclusion complexes of steroid hormones with cyclodextrins studied by the Humel–Dreyer method using reversed-phase liquid chromatography

G.S. Clarke 705 The use of temperature to empirically determine system suitability validation parameters for liquid chromatography

V. Reichelova, J. Liliemark and F. Albertioni 711 Liquid chromatographic study of acid stability of 2-chloro-2′-*arabino*-fluoro-2′-deoxyadenosine, 2-chloro-2′-deoxyadenosine and related analogues

L. Yakushina and A. Taranova 715 Rapid HPLC simultaneous determination of fat-soluble vitamins, including carotenoids, in human serum

Z. Lin and O. Beck 719 Procedure for verification of flunitrazepam and nitrazepam intake by gas chromatographic–mass spectrometric analysis of urine

Number 6

Full Papers

D.A. Ryan, J.K.M. Sanders, G.C. Curtis and H. Hughes 723 NMR study of whole rat bile: the biliary excretion of cefoperazone and benzyl chloride by an isolated perfused rat liver

D.A. Ryan, J.K.M. Sanders, G.C. Curtis and H. Hughes 735 NMR study of whole rat bile: the biliary excretion of 4-cyano-N,N-dimethyl aniline by an isolated perfused rat liver and a liver *in situ*

A.G. Hausberger and P.P. DeLuca 747 Characterization of biodegradable poly(D,L-lactide-co-glycolide) polymers and microspheres

I.I. Hewala 761 Spectrofluorimetric and derivative absorption spectrophotometric techniques for the determination of loperamide hydrochloride in pharmaceutical formulations

J.A. Murillo, J.M. Lemus and L.F. García 769 Simultaneous determination of the binary mixtures of cefsulodin and clavulanic acid by using first-derivative spectrophotometry

A.M. Wahbi, A.M. El-Walily, E.M. Hassan, F.G. Saliman and A. El-Gendi 777 Liquid chromatographic determination of flunarizine dihydrochloride in the presence of its degradation product

Short Communications

L. Edno, F. Bressolle, B. Combe and M. Galtier 785 A reproducible and rapid HPLC assay for quantitation of piroxicam in plasma

F. Kamali and E. Nicholson 791 Determination of 3,4-diaminopyridine in human plasma by high-performance liquid chromatography

795 Erratum

Number 7

Review

M.D. Luque de Castro, J.M. Fernández-Romero and R. Quiles 797 Determination of inorganic ions of clinical interest: state-of-the-art and trends

Full Papers

C.L. Flurer 809 The analysis of aminoglycoside antibiotics by capillary electrophoresis

O. Petitjean, M. Tod and K. Louchahi 817 Influence of methodological aspects on tissue drug concentration estimation

M. Lázníček and A. Lázníčková 823 The effect of lipophilicity on the protein binding and blood cell uptake of some acidic drugs

A. Marsh, M.B. Evans and J. Strang 829 Radioimmunoassay of drugs of abuse in hair. Part 2: The determination of methadone in the hair of known drug users

L. Pridal, S.H. Ingwersen, F.S. Larsen, J.J. Holst, K. Adelhorst and O. Kirk 841 Comparison of sandwich enzyme-linked immunoadsorbent assay and radioimmunoassay for determination of exogenous glucagon-like peptide-1(7–36)amide in plasma

A.F.M. El Walily, S.F. Belal, E.A. Heaba and A. El Kersh 851 Simultaneous determination of enalapril maleate and hydrochlorothiazide by first-derivative ultraviolet spectrophotometry and high-performance liquid chromatography

W.-N. Wu and M.S. Mutter 857 Biotransformation of linogliride, a hypoglycemic agent in laboratory animals and humans

S. Levin, R. Nudelman, P. Reschiglian and L. Pasti 869 Simulation of fractograms of fat emulsions in power-programmed sedimentation field-flow fractionation (SdFFF)

N. Lewen, M. Schenkenberger, T. Larkin, S. Conder and H.G. Brittain 879 The determination of palladium in fosinopril sodium (monopril) by ICP-MS

J. Kyung Kim, A. Adam, J.C.K. Loo and H. Ong 885 A chemiluminescence enzyme immunoassay (CLEIA) for the determination of medroxyprogesterone acetate in human serum

M.I. Davies, C.E. Lunte and M.R. Smyth 893 Use of micellar electrokinetic capillary chromatography in the study of in vitro metabolism of phenol by human liver microsomes

J. Sochor, J. Klimeš, J. Sedláček and M. Zahradníček 899 Determination of ibuprofen in erythrocytes and plasma by high performance liquid chromatography

W. Naidong, S. Hua, E. Roets and J. Hoogmartens 905 Assay and purity control of minocycline by thin-layer chromatography using UV and fluorescence densitometry — A comparison with liquid chromatography

D.R. Brocks, M.J. Dennis and W.H. Schaefer 911 A liquid chromatographic assay for the stereospecific quantitative analysis of halofantrine in human plasma

S. Stavchansky, S. Demirbas, L. Reyderman and C.-K. Chai 919 Simultaneous determination of dextrorphan and guaifenesin in human plasma by liquid chromatography with fluorescence detection

E. Hvattum, P.T. Normann, G.C. Jamieson, J.-J. Lai and T. Skotland — 927 Detection and quantitation of gadolinium chelates in human serum and urine by high-performance liquid chromatography and post-column derivatization of gadolinium with Arsenazo III

Short Communication
S. Vladimirov, J. Brboric, M. Švonja and D. Zivanov-Stakic — 933 Spectrophotometric determination of nizatidine in pharmaceutical formulations

Number 8

Full Papers
J.D. Gilbert, T.F. Greber, J.D. Ellis, A. Barrish, T.V. Olah, C. Fernández-Metzler, A.S. Yuan and C.J. Burke — 937 The development and cross-validation of methods based on radioimmunoassay and LC/MS-MS for the quantification of the Class III antiarrhythmic agent, MK-0499, in human plasma and urine

K.D. Altria, P. Frake, I. Gill, T. Hadgett, M.A. Kelly and D.R. Rudd — 951 Validated capillary electrophoresis method for the assay of a range of basic drugs

S.L. Tamisier-Karolak, M. Tod, P. Bonnardel, M. Czok and P. Cardot — 959 Daily validation procedure of chromatographic assay using gaussoexponential modelling

R.D. Farrant, M. Spraul, I.D. Wilson, A.W. Nicholls, J.K. Nicholson and J.C. Lindon — 971 Assignment of the 750 MHz ^{1}H NMR resonances from a mixture of transacylated ester glucuronic acid conjugates with the aid of oversampling and digital filtering during acquisition

K.L. Hoffman, D.T. Rossi and D.S. Wright — 979 Trace quantitation of the novel cholinesterase inhibitor in human plasma by capillary gas chromatography/ion trap mass spectrometry

V. Segarra, F. Carrera, J.L. Fabregas and J. Claramunt — 987 Degradation profile and identification of the major degradation products of dobupride under several conditions by GC/MS and HPLC-profile beam/MS

A. Acheampong and D.D.-S. Tang-Liu — 995 Measurement of brimonidine concentrations in human plasma by a highly sensitive gas chromatography/mass spectrometric assay

P. Umapathi and P. Parimoo — 1003 Second-order derivative spectrophotometric assay for imipramine hydrochloride and diazepam in pure admixtures and in dosage forms

M.B. Brown, J.N. Miller and N.J. Seare — 1011 An investigation of the use of Nile Red as a long-wavelength fluorescent probe for the study of α_1-acid glycoprotein–drug interactions

C. Cruces Blanco, A. Segura Carretero, A. Fernández Gutierrez and M. Román Ceba — 1019 Micellar-enhanced synchronous-derivative fluorescence determination of derivatized folic acid in pharmaceutical preparations

S.P. Singh and D.E. Moody — 1027 A radiometric TLC assay of liver microsomal dextromethorphan O-demethylation

S.S. Al-Obaidy, A. Li Wan Po, P.J. McKiernan, J.F.T Glasgow and J. Millership — 1033 Assay of paracetamol and its metabolites in urine, plasma and saliva of children with chronic liver disease

F.T. Delbeke, J. Landuyt and M. Debackere — 1041 Disposition of human drug preparations in the horse. IV. Orally administered fenoprofen

Short Communications
D.J. Kroon, J. Freedy, D.J. Burinsky and B. Sharma — 1049 Rapid profiling of carbohydrate glycoforms in monoclonal antibodies using MALDI/TOF mass spectrometry

D.G. Reed, G.P. Martin, J.M. Konieczny and M.A. Brooks — 1055 The determination of alendronate sodium in tablets by inductively coupled plasma (ICP)

M. Santos, E. García, F.G. López, J.M. Lanao and A. Domínguez-Gil — 1059 Determination of netilmicin in plasma by HPLC

Number 9

Analytical Survey

A.C. Calokerinos, N.T. Deftereos and W.R.G. Baeyens 1063 Chemiluminescence in drug assay

Full Papers

R.B. Burns, J.R. Heggie and L. Embree 1073 A gas-chromatographic assay method for busulfan with sensitivity for test dose therapeutic monitoring

R. Wong, B. Leyland-Jones and I.W. Wainer 1079 A competitive enzyme linked immunosorbent assay for the determination of *N*-acetyltransferase (NAT2) phenotypes

C. Bertucci, G. Ascoli, G. Uccello-Barretta, L. Di Bari and P. Salvadori 1087 The binding of 5-fluorouracil to native and modified human serum albumin: UV, CD, and ^{1}H and ^{19}F NMR investigation

V. Ródenas, A. Parra, J. Garcia-Villanova and M.D. Gómez 1095 Simultaneous determination of cefepime and L-arginine in injections by second-derivative spectrophotometry

T. Pérez-Ruiz, C. Martínez-Lozano, A. Sanz and S. Mondéjar 1101 Flow-injection extraction-spectrophotometric determination of bromhexine with orange IV

M. Sanchez Pena, M.J. Acedo, F. Salinas, M.C. Mahedero and J.J. Aaron 1107 Analysis of sulfamethazine in the presence of sulfamerazine or sulfadiazine by first-derivative photochemically induced fluorescence

M.I. Albero, C. Sanches-Pedreño and M.S. Garcia 1113 Flow-injection spectrofluorimetric determination of flufenamic and mefenamic acid in pharmaceuticals

J.L. Vilchez, R. Blanc, R. Avidad and A. Navalón 1119 Spectrofluorimetric determination of paracetamol in pharmaceuticals and biological fluids

F.A. Aly, F. Belal and M.I. Walash 1127 Polarographic determination of prenalterol hydrochloride through treatment with nitrous acid

P. Leroy, O. Baudrit, M.-D. Blanchin, A. Nicolas and H. Fabre 1133 Electrochemical oxidation of CBS-113 A, a new anti-inflammatory drug: applicability to liquid chromatography-electrochemical detection

F. Jorquera, M.M. Almar, A. Jimeno, M. González-Sastre and J. González-Gallego 1141 Assessment of antipyrine kinetics from saliva or plasma: influence of age

M. Pospíšilová, V. Jokl, M. Polášek and J. Mokrá 1147 Determination of some non-steroid anti-inflammatory drugs by capillary isotachophoresis

J. Paesen, W. Cypers, K. Pauwels, E. Roets and J. Hoogmartens 1153 Study of the stability of tylosin A in aqueous solutions

G.W. Ponder and J.T. Stewart 1161 A liquid chromatographic method for the determination of promethazine enantiomers in human urine and serum using solid-phase extraction and fluorescence detection

S. Torrado, S. Fraile, J.J. Torrado and E. Selles 1167 Comparison of reversed-phase liquid chromatography with colorimetry for analysis of phenolphthalein preparations

V. Augey, M. Cociglio, M. Galtier, R. Yearoo, V. Pinsani and F. Bressolle 1173 High-performance liquid chromatographic determination of *cis*-dichlorodiammineplatinum(II) in plasma ultrafiltrate

C.M. Chavez, M.L. Constanzer and B.K. Matuszewski 1179 Determination of pirenzepine in human plasma using liquid chromatography with tandem mass spectrometric detection

Short Communications

L. Ye, J.T. Stewart and H. Zhang 1185 A comparison of disc and cartridge solid-phase extraction for the LC determination of rifampin and 25-desacetylrifampin in human serum

G.A. Portmann and D.M. Simmons 1189 Microscopic determination of drug solubility in plasma and calculation of injection rates with a plasma circulatory model to prevent precipitation on intravenous injection

Number 10

Papers from the Analysis and Pharmaceutical Quality Section presented at the Ninth Annual Meeting of the American Association of Pharmaceutical Scientists, San Diego, CA, 6–10 November 1994

v Editorial

Full Papers

J.M. Schuette, U. Pieles, S.D. Maleknia, G.S. Srivatsa, D.L. Cole, H.E. Moser and N.B. Afeyan — 1195 Sequence analysis of phosphorothioate oligonucleotides via matrix-assisted laser desorption ionization time-of-flight mass spectrometry

J.R. Kagel, D.T. Rossi, G.D. Nordblom, R.C. Dudeck, C.M. Barksdale, B.-S. Kuo and D.S. Wright — 1205 Considerations in the development of a sensitive HPLC assay for human epidermal growth factors in human plasma

J. Zagrobelny, C. Chavez, M. Constanzer and B.K. Matuszewski — 1215 Determination of a glycine/NMDA receptor antagonist in human plasma and urine using column-switching high-performance liquid chromatography with ultraviolet fluorescence and tandem mass spectrometric detection

S.R. Knaub, M.J. Priston, M.D. Morton, J.D. Slechta, D.G. Vander Velde and C.M. Riley — 1225 A ^{19}F NMR study of lomefloxacin in human erythrocytes and its interaction with hemoglobin

K.P. Shah, M. Chang and C.M. Riley — 1235 Automated analytical systems for drug development studies. 3. Multivessel dissolution testing system based on microdialysis sampling

R.T. Foster, R.A. Carr, F.M. Pasutto and J.A. Longstreth — 1243 Stereospecific high-performance liquid chromatographic assay of lomefloxacin in human plasma

C. Hekman, S. Park, W.-Y. Teng, N.A. Guzman and T. Rossi — 1249 Degradation of lyophilized and reconstituted MACROSCINT® (DTPA-IgG): precipitation vs. glucosylation

G.A. Eberlein — 1263 Quantitation of proteins using HPLC-detector response rather than standard curve comparison

J.D. Kirsch and J.K. Drennen — 1273 Determination of film-coated tablet parameters by near-infrared spectroscopy

S. Indiran Pather, S.H. Neau and S. Pather — 1283 A comparison of two quality assessment methods for emulsions

Short Communication

N.K. Jagota, J.B. Nair and P.T. Kurtulik — 1291 Ion chromatography of methanesulfonic acid in pharmaceuticals

Number 11

Full Papers

F. Ortiz-Boyer, M.T. Tena, M.D. Luque de Castro and M. Valcárcel — 1297 Development and validation of chromatographic methods (HPLC and GC) for the determination of the active components (benzocaine, tyrothricin and menthol) of a pharmaceutical preparation

R. Brindle, K. Albert, E.D. Morgan, P. Martin and I.D. Wilson — 1305 Solid state NMR and extraction studies on "phenyl"-bonded stationary phases used for solid phase extraction

G.M. Hanna and C.A. Lau-Cam — 1313 Determination of the optical purity of timolol maleate by proton nuclear magnetic resonance spectroscopy with a chiral Pr(III) shift reagent

D. Bonazzi, V. Andrisano, R. Gatti and V. Cavrini — 1321 Analysis of pharmaceutical creams: a useful approach based on solid-phase extraction (SPE) and UV spectrophotometry

L.A. Carreira, M. Rizk, Y. El-Shabrawy, N.A. Zakhari and S.S. Toubar — 1331 Europium(II) ion probe spectrofluorometric determination of diclofenac sodium

D. Obendorf and G. Stubauer 1339 Adsorptive stripping voltammetry of nicardipine at a HMDE: determination of trace levels nicardipine in blood and urine

C. Bottalico, G. Micelli, A. Guerrieri, F. Palmisano, V. Lorusso and M. De Lena 1349 An on-line semi-automated solid-phase extraction procedure for high-performance liquid chromatographic determination of lonidamine in serum

K.M. Bradshaw, J. Burnett and A.S. Sidhu 1355 High-performance liquid chromatographic determination of bisacodyl in pharmaceutical dosage forms marketed in Australia

M.E.A. Faouzi, T. Dine, M. Luyckx, C. Brunet, M.-L. Mallevais, F. Goudaliez, B. Gressier, M. Cazin, J. Kablan and J.C. Cazin 1363 Stability, compatibility and plasticizer extraction of miconazole injection added to infusion solutions and stored in PVC containers

B. Voith, H. Spahn-Langguth and E. Mutschler 1373 New specific and sensitive HPLC-assays for ethacrynic acid and its main metabolite — the cysteine conjugate — in biological material

S.D. Studenberg, J.D. Long, J.H. Woolf, C.J. Bruner, D. Wilson and J.L. Woolley 1383 A robotics-based liquid chromatographic assay for the measurement of atovaquone in plasma

H. Liu, H. Davoudi and T. Last 1395 Determination of Amphotericin B in cerebrospinal fluid by solid-phase extraction and liquid chromatography

K.M. Höld, D. de Boer, J.R. Soedirman, J. Zuidema and R.A.A. Maes 1401 The secretion of propranolol enantiomers in human saliva: evidence for active transport?

Short Communications

C.-S. Liu, Y.S. Song, K.-J. Zhang, J.-C. Ryu, M. Kim and T.-H. Zhou 1409 Gas chromatographic mass spectrometric profiling of luteolin and its metabolites in rat urine and bile

H. Kim and C.-C. Lin 1415 A chiral liquid chromatographic method for the determination of the enantiomers of the racemic triazole antifungal drug (SCH 39304) in human plasma

J.A. Troskosky, T. Katona, J.P. Zodda and M.N. Eakins 1421 Determination of trace levels of Dy^{3-} in Dy(HP-DO3A) by ion-pair liquid chromatography with post-column reaction

Number 12

Full Papers

K.A. Selinger 1427 Inspection by variables as an acceptance criterion in bioanalysis — a proposal

R. Jones 1437 Quantitation of mixtures from two-dimensional data sets using orthonormal functions

S. Khundker, J.R. Dean and P. Jones 1441 A comparison between solid phase extraction and supercritical fluid extraction for the determination of fluconazole from animal feed

B. Gallez, J.-L. De Keyser, R. Debuyst, F. Dejehet, L. Neuvens and P. Dumont 1449 The effects of pasteurisation on albumin: an EPR binding assay for polymeric albumin

E. Kiliç, F. Köseoğlu, A. Kenar and M. Abdülkadir Akay 1453 Conductimetric titrimetry for assay of selected antibiotics in non-aqueous media

R.G.H. van Leuken, A.L.L. Duchateau, and G.T.C. Kwakkenbos 1459 Thermospray liquid chromatography/mass spectrometry study of diastereomeric isoindole derivatives of amino acids and amino acid amides

M. Takamoto, T. Yano, T. Shintani and S. Hiraku 1465 A highly sensitive and selective method for the determination of Leukotriene B_4 in human plasma by negative ion chemical ionization/gas chromatography/ tandem mass spectrometry

I. Bjørnsdottir and S.H. Hansen	1473	Determination of opium alkaloids in crude opium using non-aqueous capillary electrophoresis
H. Nishi, K. Ishibuchi, K. Nakamura, H. Nakai and T. Sato	1483	Enantiomeric separation of denopamine by capillary electrophoresis and high-performance liquid chromatography using cyclodextrins
G.K. Poon, P. Mistry, F.I. Raynaud, K.R. Harrap, B.A. Murrer and C.F.J. Barnard	1493	Determination of metabolites of a novel platinum anticancer drug JM216 in human plasma ultrafiltrates
G.S. Rekhi, S.S. Jambhekar, P.F. Souney and D.A. Williams	1499	A fluorimetric liquid chromatography method for the determination of propanolol in human serum/plasma
H. Tajerzadeh and S. Dadashzadeh	1507	An isocratic high-performance liquid chromatographic system for simultaneous determination of theophylline and its major metabolites in human urine
D.A. Roston, J.J. Sun, P.W. Collins, W.E. Perkins and S.J. Tremont	1513	Supercritical fluid extraction–liquid chromatography method development for a polymeric controlled-release drug formulation
K. Selinger, G. Smith, S. Depee and C. Aureche	1521	Determination of GI147211 in human blood by HPLC with fluorescence detection
M.-C. Tzou, R.A. Sams and R.H. Reuning	1531	Specific and sensitive determination of digoxin and metabolites in human serum by high performance liquid chromatography with cyclodextrin solid-phase extraction and precolumn fluorescence derivatization
P.A. Coates, I.S. Blagbrough, T. Lewis, B.V.L. Potter and M.G. Rowan	1541	An HPLC assay for the norditerpenoid alkaloid methyllycaconitine, a potent nicotinic acetylcholine receptor antagonist
R.W. Sparidans, J. den Hartigh and P. Vermeij	1545	High-performance ion-exchange chromatography with in-line complexation of bisphosphonates and their quality control in pharmaceutical preparations
Short Communications		
L.B. Pfendt, G.V. Popović and S.I. Supančić	1551	Spectrophotometric determination of midazolam in pharmaceutical formulations
G. Indrayanto, A. Sunarto and Y. Adriani	1555	Simultaneous assay of phenylpropanolamine hydrochloride, caffeine, paracetamol, glycerylguaiacolate and chlorpheniramine maleate in Silabat™ tablet using HPLC with diode array detection
Book Reviews	1561	
Volume Contents, Author Index and Subject Index, Volume 13, 1995	iii	